

Algorithms for Determining Birationality of Parametrization of Affine Curves

Hyungju Park

Department of Mathematics and Statistics

Oakland University

Rochester, MI 48309

email: park@oakland.edu

Abstract

Let k be an arbitrary field, and C be a curve in $\mathbb{A}^n$ defined parametrically by $x_1 = f_1(t), \dots, x_n = f_n(t)$, where $f_1, \dots, f_n \in k[t]$. A necessary and sufficient condition for the two function fields $k(t)$ and $k(f_1, \dots, f_n)$ to be same is developed in terms of zero-dimensionality of a derived ideal in the bivariate polynomial ring $k[s, t]$. Since zero-dimensionality of such an ideal can be readily determined by a Gröbner basis computation, this gives an algorithm that determines if the parametrization $\psi = (f_1, \dots, f_n) : \mathbb{A} \rightarrow C$ is a birational equivalence. We also develop an algorithm that determines if $k[t]$ and $k[f_1, \dots, f_n]$ are same, by which we get an algorithm that determines if the parametrization $\psi = (f_1, \dots, f_n) : \mathbb{A} \rightarrow C$ is an isomorphism. We include some computational examples showing the application of these algorithms.

1991 Mathematics Subject Classification: 13P10

1 Introduction

For an arbitrary field k , let $f_1, \dots, f_n \in k[t]$. Consider the curve $C \subset \mathbb{A}_k^n$ given parametrically by

$$x_1 = f_1(t), \dots, x_n = f_n(t).$$

This paper is concerned with the following two questions on this parametrization of C :

Q1. Can we effectively determine if the parametrization

$$\psi = (f_1, \dots, f_n) : \mathbb{A} \rightarrow C$$

is a birational equivalence, or equivalently $k(f_1, \dots, f_n) = k(t)$?

Q2. If the answer to Q1 is yes, can we determine if the parametrization ψ is an isomorphism, or equivalently $k[f_1, \dots, f_n] = k[t]$?

For the case of two polynomials $f_1, f_2 \in k[t]$, there are several results available. S. Abhyankar and T. Moh [AM75] have given a necessary condition for $k[f_1, f_2] = k[t]$ in terms of degrees of f_1 and f_2 :

Theorem 1.1 (S. Abhyankar and T. Moh, [AM75]) *Let k be an arbitrary field of characteristic p ($p = 0$ or $p > 0$). Suppose that f_1 and f_2 are in $k[t]$ with $m := \deg(f_1) \leq n := \deg(f_2)$, and that p does not divide $\gcd(m, n)$. If $k[f_1, f_2] = k[t]$, then m divides n .*

A. van den Essen and J. Yu [vdEY97] introduced the notion of D -resultant $D(s) \in k[s]$ of f and g , and gave necessary and sufficient conditions for $k(f, g) = k(t)$ and $k[f, g] = k[t]$ in terms of $D(s)$ (Theorem 2.1, [vdEY97]).

Example 1.2 Consider $f_1(t) = t^3, f_2(t) = t^2 + t \in k[t]$. The function field $k(t)$ is an algebraic extension of its subfield $k(t^3, t^2 + t)$. To determine if $k(t^3, t^2 + t) = k(t)$, denote the field $k(t^3, t^2 + t)$ by K . Then

$$\begin{aligned} t^2 + t \in K &\implies (t^2 + t)^2 = t^4 + 2t^3 + t^2 = t^4 - t + (t^2 + t) + 2t^3 \in K \\ &\implies t^4 - t = t(t^3 - 1) \in K \quad (\text{Since } t^2 + t, 2t^3 \in K) \\ &\implies t \in K \quad (\text{Since } t^3 - 1 \in K). \end{aligned}$$

This shows $k(t^3, t^2 + t) = k(t)$. However, since the Abhyankar-Moh necessary condition is not satisfied, $k[f_1, f_2] \subsetneq k[t]$. This means that the parametrization $\psi = (t^3, t^2 + t) : \mathbb{A} \rightarrow C$ is a birational equivalence, but not an isomorphism. $\square$

In this paper, we give algorithmic solutions to Q1 and Q2 in the general case of n polynomials using the method of Gröbner bases.

2 Birational and Isomorphic Parametrization of Curves

Throughout this paper, unless there is a possibility of confusion, we will use the shorthand notation $\mathbb{A}^n$ for the affine space $\mathbb{A}_k^n$.

For an arbitrary field k , consider the curve $C \subset \mathbb{A}^n$ given parametrically by

$$x_1 = f_1(t), \dots, x_n = f_n(t),$$

where $f_1, \dots, f_n \in k[t]$. The morphism ψ defined by

$$\psi = (f_1, \dots, f_n) : \mathbb{A} \rightarrow \mathbb{A}^n,$$

will be simply referred to as the parametrization of C by polynomials $f_1, \dots, f_n$. The curve C is the Zariski closure $\overline{\text{Im}(\psi)}$ of $\text{Im}(\psi)$ in $\mathbb{A}^n$, and we occasionally identify the morphism $\psi : \mathbb{A} \rightarrow \mathbb{A}^n$ with the induced morphism $\psi : \mathbb{A} \rightarrow C$,

Problem Q1 is equivalent to determining if the induced map of the functions fields

$$\begin{aligned} \psi^* : K(\mathbb{A}^n) = k(x_1, \dots, x_n) &\longrightarrow K(\mathbb{A}) = k(t) \\ x_i &\longmapsto f_i. \end{aligned}$$

is surjective, i.e.

$$k(f_1, \dots, f_n) = k(t).$$

So we need to understand what conditions on the polynomials $f_1, \dots, f_n$ will guarantee $k(f_1, \dots, f_n) = k(t)$.

For Problem Q2, consider the induced k -algebra homomorphism ψ^* of the coordinate rings

$$\begin{aligned} \psi^* : A(\mathbb{A}^n) = k[x_1, \dots, x_n] &\longrightarrow A(\mathbb{A}) = k[t] \\ x_i &\longmapsto f_i. \end{aligned}$$

Note that the coordinate ring of $C = \overline{\text{Im}(\psi)}$ is

$$A(C) = k[x_1, \dots, x_n] / \text{Ker}(\psi^*) = \text{Im}(\psi^*) \cong k[f_1, \dots, f_n].$$

Proving that ψ is an immersion is equivalent to proving that their coordinate rings are isomorphic:

$$k[t] = k[f_1, \dots, f_n].$$

Now we will describe the injectivity and birationality of

$$\psi : \mathbb{A} \rightarrow C \subset \mathbb{A}^n$$

in terms of a set of bivariate polynomials derived from f_i 's.

Lemma 2.1 *Suppose that $f'_1, \dots, f'_n \in k[t]$ are not identically zero. Then, the morphism*

$$\psi := (f_1, \dots, f_n) : \mathbb{A} \rightarrow \mathbb{A}^n$$

is finite.

Proof: We may assume, without loss of generality, that f_1 is not a constant. Consider

$$\begin{aligned} \psi^* : k[x_1, \dots, x_n] &\longrightarrow k[t] \\ x_i &\longmapsto f_i. \end{aligned}$$

Dividing f_1 by its leading coefficient if necessary, we may assume $f_1(t)$ is monic. Then the monic polynomial

$$G(T) := f_1(T) - f_1 \in k[f_1, \dots, f_n](T)$$

gives an integral dependence of $t \in k[t]$ on $k[f_1, \dots, f_n]$. $\square$

For $f_1, \dots, f_n \in k[t]$, we introduce a new variable s and consider

$$f_1(t), \dots, f_n(t), f_1(s), \dots, f_n(s) \in k[s, t].$$

For each $i = 1, \dots, n$, $t-s$ divides $f_i(t) - f_i(s)$ and there exists $g_i(s, t) \in k[s, t]$ such that $f_i(t) - f_i(s) = (t-s)g_i(s, t)$. We will identify the fraction $\frac{f_i(t) - f_i(s)}{t-s}$ with the polynomial $g_i(s, t) \in k[s, t]$. It is easy to prove that $g_i(s, s) = f'_i(s)$. The following theorem characterizes the algebraic set $V(g_1, \dots, g_n) \subset \mathbb{A}^2$.

Theorem 2.2 *For $f_1, \dots, f_n \in k[t]$, let*

$$g_i(s, t) := \frac{f_i(t) - f_i(s)}{t-s} \in k[s, t], \quad i = 1, \dots, n.$$

Then, for the morphism $\psi := (f_1, \dots, f_n) : \mathbb{A} \rightarrow \mathbb{A}^n$,

$$V(g_1, \dots, g_n) = A_\psi \amalg B_\psi, \tag{2.1}$$

where

$$\begin{aligned} A_\psi &= \{(a, b) \mid a \neq b \in k \text{ and } \psi(a) = \psi(b)\} \\ B_\psi &= \{(a, a) \mid a \in V(f'_1, \dots, f'_n)\}. \end{aligned}$$

Proof: Suppose $(a, b) \in A_\psi$. Then,

$$\begin{aligned}
(a, b) \in A_\psi &\implies a \neq b \text{ and } \psi(a) = \psi(b) \\
&\implies a \neq b \text{ and } f_i(a) = f_i(b), \forall i = 1, \dots, n \\
&\implies g_i(a, b) = 0, \forall i = 1, \dots, n \\
&\implies (a, b) \in V(g_1, \dots, g_n).
\end{aligned}$$

Hence, $A_\psi \subset V(g_1, \dots, g_n)$.

Suppose $(a, a) \in B_\psi$. Then,

$$\begin{aligned}
(a, a) \in B_\psi &\implies f'_i(a) = 0, \forall i = 1, \dots, n \\
&\implies g_i(a, a) = 0, \forall i = 1, \dots, n \\
&\implies (a, a) \in V(g_1, \dots, g_n).
\end{aligned}$$

Hence $B_\psi \subset V(g_1, \dots, g_n)$. Therefore, $A_\psi \amalg B_\psi \subset V(g_1, \dots, g_n)$.

In order to show $V(g_1, \dots, g_n) \subset A_\psi \amalg B_\psi$, let $(a, b) \in V(g_1, \dots, g_n)$.

If $a \neq b$, then

$$\begin{aligned}
g_i(a, b) &= \frac{f_i(b) - f_i(a)}{b - a} = 0, \forall i = 1, \dots, n \\
&\implies f_i(b) = f_i(a), \forall i = 1, \dots, n \\
&\implies \psi(a) = \psi(b) \\
&\implies (a, b) \in A_\psi.
\end{aligned}$$

If $a = b$, then

$$\begin{aligned}
g_i(a, a) &= 0, \forall i = 1, \dots, n \\
&\implies a \in V(g_1(s, s), \dots, g_n(s, s)) = V(f'_1(s), \dots, f'_n(s)) \\
&\implies (a, a) \in B_\psi.
\end{aligned}$$

□

Remark 2.3 The set A_ψ in Theorem 2.2 describes the multiple points on the curve C , while the set B_ψ describes the ramification points (or branch points) on C . More precisely, if $(a, b) \in A_\psi$, then the point $\psi(a) = \psi(b)$ on C has multiplicity of at least 2. If $(a, a) \in B_\psi$, then $\psi(a)$ is a ramification point on C . Therefore, if $B_\psi = \emptyset$, then the parametrization $\psi : \mathbb{A} \rightarrow C$ is an étale morphism. □

Theorem 2.4 For $f_1, \dots, f_n \in k[t]$, let

$$g_i(s, t) := \frac{f_i(t) - f_i(s)}{t - s} \in k[s, t], \quad i = 1, \dots, n.$$

Suppose that $f'_1, \dots, f'_n \in k[t]$ are not identically zero. Then, the morphism

$$\psi := (f_1, \dots, f_n) : \mathbb{A} \rightarrow \mathbb{A}^n$$

is a closed immersion if and only if $V(g_1, \dots, g_n) = \emptyset$.

Proof: Suppose that $V(g_1, \dots, g_n) = \emptyset$ and C is the curve in $\mathbb{A}^n$ given parametrically by $x_1 = f_1(t), \dots, x_n = f_n(t)$. By Lemma 2.1, $\psi : \mathbb{A} \rightarrow C$ is finite, and thus proper. Since

$$A_\psi = B_\psi = \emptyset,$$

by Remark 2.3, C is nonsingular, and thus normal. Since each fiber of the finite morphism $\psi : \mathbb{A} \rightarrow C$ contains one point, its degree $[K(\mathbb{A}) : K(C)]$ is 1, that is,

$$k(f_1, \dots, f_n) = k(t).$$

The normality of C implies that

$$A(C) = k[f_1, \dots, f_n] \subset k[t]$$

is integrally closed in $K(C) = k(f_1, \dots, f_n) = k(t)$. Since the normal ring $k[t]$ is integral over $k[f_1, \dots, f_n]$, we conclude

$$k[f_1, \dots, f_n] = k[t].$$

Therefore, the morphism $\psi : \mathbb{A} \rightarrow \mathbb{A}^n$ induces an isomorphism of $\mathbb{A}$ onto the closed set C in $\mathbb{A}^n$.

Conversely, if $\psi : \mathbb{A} \rightarrow \mathbb{A}^n$ is a closed immersion, then $C \cong \mathbb{A}$ is nonsingular. By Remark 2.3,

$$A_\psi = B_\psi = \emptyset.$$

Hence $V(g_1, \dots, g_n) = \emptyset$. □

Corollary 2.5 For $f_1, \dots, f_n \in k[t]$, let

$$g_i(s, t) := \frac{f_i(t) - f_i(s)}{t - s} \in k[s, t], \quad i = 1, \dots, n.$$

Suppose that $f'_1, \dots, f'_n \in k[t]$ are not identically zero. Then,

$$k[f_1, \dots, f_n] = k[t]$$

if and only if $V(g_1, \dots, g_n) = \emptyset$.

Theorem 2.6 For $f_1, \dots, f_n \in k[t]$, let

$$g_i(s, t) := \frac{f_i(t) - f_i(s)}{t - s} \in k[s, t], \quad i = 1, \dots, n.$$

Suppose that $f'_1, \dots, f'_n \in k[t]$ are not identically zero and C is the curve in $\mathbb{A}^n$ given parametrically by $x_1 = f_1(t), \dots, x_n = f_n(t)$. Then the parametrization

$$\psi = (f_1, \dots, f_n) : \mathbb{A} \rightarrow C$$

is a birational equivalence if and only if $V(g_1, \dots, g_n)$ is a finite set.

Proof: By Lemma 2.1, the morphism $\psi : \mathbb{A} \rightarrow \mathbb{A}^n$ is finite, and thus proper. Hence $\text{Im}(\psi)$ is a Zariski closed set of dimension 1 in $\mathbb{A}^n$, and is equal to C .

($\Leftarrow$) Let $V(g_1, \dots, g_n) = \{(a_1, b_1), \dots, (a_l, b_l)\}$. Define open sets $U \subset \mathbb{A}$ and $V \subset \mathbb{A}^n$ by

$$U = \mathbb{A} - \{a_1, \dots, a_l\}, \quad V = C - \{\psi(a_1), \dots, \psi(a_l)\}.$$

Then ψ induces an finite injective morphism $\psi|_U : U \rightarrow V$. Since each fiber of $\psi|_U$ has one point, its degree $[K(U) : K(V)]$ is 1. Therefore, $\psi : \mathbb{A} \rightarrow C$ is birational.

($\Rightarrow$) Since $\psi : \mathbb{A} \rightarrow C = \text{Im}(\psi)$ is birational, there exist open sets $U \subset \mathbb{A}$ and $V \subset C$ such that ψ induces an isomorphism between them. Since $\mathbb{A} - U$ is a proper closed subset of $\mathbb{A}$, the irreducibility of $\mathbb{A}$ forces $\dim(\mathbb{A} - U) < 1$, i.e. $\mathbb{A} - U$ is a finite set. Therefore, the injectivity of ψ fails only at finitely many points of $\mathbb{A}$, and A_ψ is a finite set. Since at least one of f'_i 's is nonzero, $f'_1, \dots, f'_n \in k[t]$ have at most finitely many zeros. This means that B_ψ is a finite set. Hence, by Theorem 2.2 $V(g_1, \dots, g_n) = A_\psi \amalg B_\psi$ is a finite set. $\square$

Corollary 2.7 For $f_1, \dots, f_n \in k[t]$, let

$$g_i(s, t) := \frac{f_i(t) - f_i(s)}{t - s} \in k[s, t], \quad i = 1, \dots, n.$$

Suppose that $f'_1, \dots, f'_n \in k[t]$ are not identically zero. Then,

$$k(f_1, \dots, f_n) = k(t)$$

if and only if $|V(g_1, \dots, g_n)| < \infty$.

Example 2.8 Consider $f_1(t) = t^3$ and $f_2(t) = t^2 + t \in k[t]$ of Example 1.2. Let us compute

$$V\left(\frac{f_1(t) - f_1(s)}{t - s}, \frac{f_2(t) - f_2(s)}{t - s}\right).$$

We have to solve

$$\begin{aligned} \frac{f_1(t) - f_1(s)}{t - s} &= \frac{t^3 - s^3}{t - s} = t^2 + ts + s^2 = 0 \\ \frac{f_2(t) - f_2(s)}{t - s} &= \frac{(t^2 - s^2) + (t - s)}{t - s} = t + s + 1 = 0. \end{aligned}$$

From the second equation, $t = -s - 1$. By putting it into the first equation, we get

$$(-s - 1)^2 + (-s - 1)s + s^2 = s^2 + s + 1 = 0.$$

If $k = \mathbb{C}$, then

$$V\left(\frac{f_1(t) - f_1(s)}{t - s}, \frac{f_2(t) - f_2(s)}{t - s}\right) = \left\{\left(\frac{-1+i\sqrt{3}}{2}, \frac{-1-i\sqrt{3}}{2}\right), \left(\frac{-1-i\sqrt{3}}{2}, \frac{-1+i\sqrt{3}}{2}\right)\right\}.$$

It can be easily seen that $V\left(\frac{f_1(t)-f_1(s)}{t-s}, \frac{f_2(t)-f_2(s)}{t-s}\right)$ is finite over an arbitrary field k . Since this set is finite, Theorem 2.6 confirms our earlier finding $k(t^3, t^2 + t) = k(t)$. But since this set is nonempty, by Theorem 2.4, we conclude that $\psi = (f, g) : \mathbb{A} \rightarrow \mathbb{A}^2$ is not a closed immersion, i.e. $k[t^3, t^2 + t] \subsetneq k[t]$ as predicted by the Abhyankar-Moh result [AM75]. $\square$

3 Algorithms

For $f_1, \dots, f_n \in k[s, t]$, define $g_1, \dots, g_n$ by $g_i(s, t) := \frac{f_i(t) - f_i(s)}{t - s}$, $i = 1, \dots, n$. The finiteness condition on $V(g_1, \dots, g_n)$ in Theorem 2.6 is equivalent to the zero-dimensionality of the ideal $I := \langle g_1, \dots, g_n \rangle \subset k[s, t]$. Fix a term ordering $\prec$ on the set of monomials in $k[s, t]$, and let $h_1, \dots, h_l \in k[s, t]$ be the reduced Gröbner basis of the ideal I w.r.t $\prec$. For an arbitrary polynomial

$f \in k[s, t]$, denote the initial (or leading) term of f w.r.t. $\prec$ by $\text{in}(f)$. Then I is zero dimensional if and only if there exist $i, j \in \{1, \dots, l\}$ such that $\text{in}(h_i) = s^p$ and $\text{in}(h_j) = t^q$ for some $p, q \in \mathbb{N}$ (see [AL94, Theorem 2.2.7] for a proof). This produces the following algorithmic solution to Problem Q1:

Algorithm 1

Input: $f_1, \dots, f_n \in k[t]$.
Output: yes if $k(f_1, \dots, f_n) = k(t)$, no otherwise.
Step 1: For each $i = 1, \dots, n$, compute $g_i := \frac{f_i(t) - f_i(s)}{t - s} \in k[s, t]$.
Step 2: Compute the reduced Gröbner basis $G = \{h_1, \dots, h_l\}$ of the ideal $I := \langle g_1, \dots, g_n \rangle \subset k[s, t]$.
Step 3: Output yes if there exist $i, j \in \{1, \dots, l\}$ such that $\text{in}(h_i) = s^p$ and $\text{in}(h_j) = t^q$ for some $p, q \in \mathbb{N}$.
Output no otherwise.

Suppose that the ground field k is algebraically closed and $g_1, \dots, g_n \in k[s, t]$. Then, by Hilbert Nullstellensatz, $V(g_1, \dots, g_n) \subset \mathbb{A}^2$ is empty if and only if $g_1, \dots, g_n \in k[s, t]$ generate the unit ideal, i.e. there exist $h_1, \dots, h_n \in k[s, t]$ such that

$$h_1 g_1 + \dots + h_n g_n = 1.$$

This observation together with Theorem 2.4 produces the following algorithmic solution to Problem Q2:

Algorithm 2 (over an algebraically closed field k)

Input: $f_1, \dots, f_n \in k[t]$.
Output: yes if $k[f_1, \dots, f_n] = k[t]$, no otherwise.
Step 1: For each $i = 1, \dots, n$, compute $g_i := \frac{f_i(t) - f_i(s)}{t - s} \in k[s, t]$.
Step 2: Compute the reduced Gröbner basis G of the ideal $I := \langle g_1, \dots, g_n \rangle \subset k[s, t]$.
Step 3: Output yes if $G = \{1\}$. Output no otherwise.

4 Examples

Most of the examples in this section are worked out with the computer algebra system *Singular* [GPS97].

Example 4.1 Consider the trivial case of the twisted cubic $C \subset \mathbb{A}^3$ given parametrically by

$$x = t, y = t^2, z = t^3,$$

which is apparently isomorphic to the affine line $\mathbb{A}$. Let $f_1 = t, f_2 = t^2, f_3 = t^3$, and $g_i(s, t) := \frac{f_i(t) - f_i(s)}{t - s}$ for each $i = 1, 2, 3$. Since

$$g_1(s, t) = \frac{f_1(t) - f_1(s)}{t - s} = 1,$$

the reduced Gröbner basis of $\{g_1, g_2, g_3\}$ w.r.t an arbitrary term ordering is $\{1\}$. Therefore, Algorithm 2 confirms that the parametrization

$$\psi = (t, t^2, t^3) : \mathbb{A} \rightarrow C$$

is an isomorphism. □

Example 4.2 Consider the curve $C \subset \mathbb{A}^2$ given parametrically by

$$x = f_1(t) := 2t^8 + t^4 + 3t + 1, \quad y = f_2(t) := t^4 - 2t^2 + 2.$$

Then,

$$\begin{aligned} g_1(s, t) &:= \frac{f_1(t) - f_1(s)}{t - s} \\ &= 2(t^7 + t^6s + t^5s^2 + t^4s^3 + t^3s^4 + t^2s^5 + ts^6 + s^7) \\ &\quad + (t^3 + t^2s + ts^2 + s^3) + 3, \\ g_2(s, t) &:= \frac{f_2(t) - f_2(s)}{t - s} \\ &= (t^3 + t^2s + ts^2 + s^3) - 2(t + s). \end{aligned}$$

Fix the degree reverse lex order $\prec$ on $k[s, t]$ with $s \prec t$. Then a computation with *Singular* shows that the reduced Gröbner basis of $\{g_1, g_2\}$ w.r.t. $\prec$ is $\{h_1, h_2, h_3, h_4\}$ where

$$\begin{aligned} h_1 &= t^2s + s^3 - 2s, \\ h_2 &= t^3 + t^2s + ts^2 + s^3 - 2t - 2s, \\ h_3 &= 8ts^4 + 8s^5 - 16ts^2 - 16s^3 + 18, \\ h_4 &= 16s^6 - 48s^4 - 18t^2 - ts + 51s^2. \end{aligned}$$

Therefore, Algorithm 1 and Algorithm 2 show that the parametrization $\psi = (f_1, f_2) : \mathbb{A} \rightarrow C$ is a birational equivalence, but not an isomorphism (over an algebraically closed field). One notes that, although $k[f_1, f_2] \neq k[t]$, the Abhyankar-Moh necessary condition [AM75] for $k[f_1, f_2] = k[t]$ is satisfied since $\deg(f_1)$ divides $\deg(f_2)$. Hence this example confirms that the Abhyankar-Moh condition is a necessary but not a sufficient condition for $k[f_1, f_2] = k[t]$. $\square$

Example 4.3 Consider the curve $C \subset \mathbb{A}^3$ given parametrically by

$$x = f_1(t) := t^{10} + t^4, \quad y = f_2(t) := t^8 + 2t^2, \quad z = f_3(t) := t^6 - t^4 + 1.$$

Then, for $g_i(s, t) := \frac{f_i(t) - f_i(s)}{t - s}$, $i = 1, 2, 3$, the reduced Gröbner basis G of $\{g_1, g_2, g_3\}$ w.r.t. the degree reverse lex order is

$$G = \{t + s\}.$$

Hence, according to Algorithm 1, the parametrization

$$\psi = (f_1, f_2, f_3) : \mathbb{A} \rightarrow C$$

is not a birational equivalence. $\square$

References

- [AL94] W. Adams and P. Lounstaunau. *An Introduction to Gröbner Bases*, volume 3 of *Graduate Studies in Mathematics*. American Mathematical Society, 1994.
- [AM75] A. Abhyankar and T. Moh. Embeddings of the line in the plane. *J. Reine Angew. Math.*, 276:148–166, 1975.
- [GPS97] G.-M. Greuel, G. Pfister, and H. Schönemann. Singular Reference Manual. In *Reports On Computer Algebra*, number 12. Centre for Computer Algebra, University of Kaiserslautern, May 1997. <http://www.mathematik.uni-kl.de/~zca/Singular>
- [Har77] R. Hartshorne. *Algebraic Geometry*, volume 52 of *Graduate Texts in Mathematics*. Springer-Verlag, 1977.

- [vdE95] A. van den Essen. Seven lectures on polynomial automorphisms. In A. van den Essen, editor, *Automorphisms of Affine Spaces*, pages 3–39. Kluwer Academic Publishers, 1995.
- [vdEY97] A. van den Essen and J. Yu. The d -resultant, singularities and the degree of unfaithfulness. *Proc. Amer. Math. Soc.*, 125(3):689–695, 1997.